



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Two Sharp-tailed Sparrows I took near Tarpon Springs, Florida, in the early spring of 1897, are referable to the new race.

In general plumage *A. c. diversus* has the richest coloring of any of the group, including even spring specimens of *A. nelsoni*. Typical examples of this form in winter resemble very closely some specimens of *nelsoni* in the coloring of the upper parts — much more than they resemble typical examples of *A. caudacutus* — but may be easily distinguished from the former at any season by their larger size, longer bill and very conspicuous dark stripes on the breast and flanks.

In measurement *diversus* and *caudacutus* are practically the same, and intermediates in coloring occurred on Pea Island in winter.

GENERAL NOTES.

Holbøll's Grebe on San Francisco Bay. — On November 30, 1900, three Grebes, which were strange to me, were seen swimming near the shore of the Presidio, one of which I shot. The bird proved to be *Colymbus holbøllii* in adult plumage and a female. No others have been seen by me since this date, although a great part of my time is spent upon the lower bay. The only other specimen I know of is a male, No. 1867 of the collection of the California Academy of Sciences, taken December 25, 1882, at Oakland. The Academy also has a single specimen taken at Monterey on December 31, 1894, which is labeled a female and is No. 1433, a record of which is found in 'California Water Birds,' No. II, on page 14. Mr. Leverett M. Loomis, in his extensive work off Monterey, has found this bird very uncommon. It seems to be very rare on lower San Francisco Bay, that is, that portion bounded by the ocean and where the bay branches to spread north and south. It is possible, but not probable, that it occurs on other portions of the bay. — WM. H. KOBBE, *Fort Mason, San Francisco, Cal.*

The Question of the Generic Name *Gavia*. — In the September, 1900, number of the 'Ornithologische Monatsberichte' (Vol. VIII, page 135), Dr. Anton Reichenow claims that the name *Gavia* Forster is a synonym of *Urinator* Cuvier, basing his assumption on the use of the name *Gavia* for a species of Gull by S. G. Gmelin in his 'Reise durch Russland' (Vol. I, page 152).